

Gene Section

Short Communication

CBLc (Cas-Br-M (murine) ecotropic retroviral transforming sequence c)

Olivier Rosnet

Centre d'Immunologie INSERM-CNRS de Marseille-Luminy Case 906, 13288 Marseille Cedex 9, France (OR)

Published in Atlas Database: September 1999

Online updated version : <http://AtlasGeneticsOncology.org/Genes/CBLcID194.html>
DOI: 10.4267/2042/37530

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 1999 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

Identity

Other names: CBL-3

HGNC (Hugo): CBLC

Location : 19q13.2

DNA/RNA

Transcription

2.8-3 kb mRNA; 1422 bp open reading frame.

Protein

Description

474 amino acids; 52.5 kDa; the CBLc protein contains the evolutionarily conserved aminoterminal phosphotyrosine binding region and the Ring finger with a Cys3HisCys4 motif found in CBL and CBLb. The phosphotyrosine binding domain is composed of three interacting domains: a four-helix bundle (4H), an EF-hand calcium binding domain, and a divergent SH2 domain; the three domains together form an integrated phosphoprotein-recognition module; the CBLc protein lacks the extensive proline-rich domain and the leucine zipper found in CBL and CBLb; only one proline-rich motif is conserved at the carboxy terminus of CBLc; thus, CBLc structurally most resembles the *C.elegans* and *D.melanogaster* CBL-related proteins, SLI-1 and D-CBL, respectively; an alternatively spliced mRNA codes form a CBLc isoform lacking a critical region in the phosphotyrosine binding domain.

Expression

The CBLc gene is ubiquitously expressed with highest expression in the aerodigestive tract (stomach, liver, pancreas, small intestine, colon, trachea, and lung), prostate, thyroid gland, adrenal gland, and salivary gland.

Localisation

Cytoplasmic.

Function

Regulation of signal transduction pathways; CBLc is recruited to the EGF (epidermal growth factor) receptor (EGFR) upon EGF stimulation and inhibits EGF stimulated MAP kinase activation.

Mutations

Note

No genomic alterations described to date.

References

Ollendorff V, Mattei M, Fournier E, Adelaide J, Lopez M, Rosnet O, Birnbaum D. A third human CBL gene is on chromosome 19. *Int J Oncol.* 1998 Dec;13(6):1159-61

Keane MM, Ettenberg SA, Nau MM, Banerjee P, Cuellar M, Penninger J, Lipkowitz S. cbl-3: a new mammalian cbl family protein. *Oncogene.* 1999 Jun 3;18(22):3365-75

This article should be referenced as such:

Rosnet O. CBLc (Cas-Br-M (murine) ecotropic retroviral transforming sequence c). *Atlas Genet Cytogenet Oncol Haematol.* 1999; 3(3):134.